

A not metrizable space filled with any n mutually disjoint self-similar spaces

Akihiko Kitada¹, Shousuke Ohmori², Tomoyuki Yamamoto^{1,2,*}

¹*Institute of Condensed-Matter Science, Comprehensive Resaerch Organization, Waseda University, 3-4-1 Okubo, Shinjuku-ku, Tokyo 169-8555, Japan*

²*Faculty of Science and Engineering, Waseda University, 3-4-1 Okubo, Shinjuku-ku, Tokyo 169-8555, Japan*

*corresponding author: tymmt@waseda.jp

Abstract

A $\Lambda(\text{Card}\Lambda \succ \aleph)$ -product space of $\{0, 1\}$ has a partition $\{X_1, \dots, X_n\}$ for any n a decomposition space of each X_i of which is self-similar. (February 16, 2015)

Keywords : self-similar, decomposition space, not metrizable space

1 Introduction

Let $(X, \tau) = (\{0, 1\}^\Lambda, \tau_0^\Lambda)$, $\text{Card}\Lambda \succ \aleph$ (aleph) be the Λ -product space of $(\{0, 1\}, \tau_0)$ where τ_0 is a discrete topology for $\{0, 1\}$. The topological space (X, τ) need not to be metrizable, then, the relation $\text{Card}\Lambda \succ \aleph$ (aleph) holds and since the space X is not metrizable, X is no more the so-called Cantor set. In the present report, we will mathematically confirm the existence of a partition $\{X_1, \dots, X_n\}$ for any n of X (**2-a**), $X_i \in (\tau \cap \mathfrak{S})^1 - \{\phi\}$, a decomposition space (i.e, a space of equivalence classes) of each element $X_i = (X_i, \tau_{X_i})^2$ of which is self-similar. Namely, all points in X_i are classified into equivalence classes and then the equivalence classes form a self-similar structure. Each equivalence class is a point of the decomposition space of X_i .

Some easily verified and well known statements are summarized in the next section for the preliminaries.

2 Preliminaries

- 2-a)** A family $\{X_1, \dots, X_n\}$ of subspace X_i of X is a partition of X provided that $X_i \cap X_j = \phi, i \neq j$ and $X_1 \cup \dots \cup X_n = X$.
- 2-b)** Any zero-dimensional (0-dim), perfect, T_0 (necessarily T_2)-space (X, τ) has a partition $\{X_1, \dots, X_n\}, X_i \in (\tau \cap \mathfrak{S}) - \{\phi\}$ for any n . Each subspace (X_i, τ_{X_i}) is a 0-dim, perfect, T_2 -space.
- 2-c)** Let X be a 0-dim, perfect, compact T_2 -space. Then, for *any* compact metric space Y , there exists a continuous map f from X onto Y .
- 2-d)** If $f : (X, \tau) \rightarrow (Y, \tau')$ is a quotient map, then the map $h : (Y, \tau') \rightarrow (\mathcal{D}_f, \tau(\mathcal{D}_f)), y \mapsto f^{-1}(y)$ is a homeomorphism. Here, the decomposition $\mathcal{D}_f$ of X and the decomposition topology $\tau(\mathcal{D}_f)$ are given by $\mathcal{D}_f = \{f^{-1}(y) \subset X; y \in Y\}$ and $\tau(\mathcal{D}_f) = \{\mathcal{U} \subset \mathcal{D}_f; \bigcup \mathcal{U} \in \tau\}$, respectively. $\mathcal{D}_f$ forms a partition of X .
- 2-e)** A space which is homeomorphic to a self-similar space is self-similar.
- 2-f)** Let (X, τ) be a 0-dim, perfect, compact, not-metrizable T_2 -space. From the Urysohn's metrization theorem (X, τ) is not second countable. In the partition $\{(X_1, \tau_{X_1}), \dots, (X_n, \tau_{X_n})\}$ of (X, τ) , there exists a number i_0 such that $(X_{i_0}, \tau_{X_{i_0}})$ is not second countable. The subspace $(X_{i_0}, \tau_{X_{i_0}})$ is a 0-dim, perfect, compact not-metrizable T_2 -space.

3 A partition $\{X_1, \dots, X_n\}$ of $(\{0, 1\}^\Lambda, \tau_0^\Lambda)$

Since $(X, \tau) = (\{0, 1\}^\Lambda, \tau_0^\Lambda), \tau_0 = 2^{\{0, 1\}}, \text{Card}\Lambda \succ \aleph$ is easily verified to be a 0-dim, perfect, compact T_2 -space, from **2-b**) there exists a partition $\{(X_1, \tau_{X_1}), \dots, (X_n, \tau_{X_n})\}$ of (X, τ) where each subspace

¹The notaion $\tau \cap \mathfrak{S}$ denotes the set of all closed and open sets (clopen sets) of (X, τ) .

² $\tau_A = \{u \cap A; u \in \tau\}$ for $A \subset X$.




Figure 1: A schematic explanation of the mathematical procedure. $\{X_1, \dots, X_n\}$ is a partition of X . A decomposition space $\mathcal{D}_{f_i} = \{f_i^{-1}(y) \subset X_i; y \in Y\}$ of subspace X_i of X generated from a self-similar space Y is self-similar. Each $f_i^{-1}(y)$ is a point of a self-similar space $\mathcal{D}_{f_i}$.

(X_i, τ_{X_i}) is a 0-dim, perfect, compact T_2 -space.³ From **2-c**), there exists a continuous map f_i from (X_i, τ_{X_i}) onto any compact, self-similar metric space (Y, τ_d) . Since (X_i, τ_{X_i}) is a compact space and Y is a T_2 -space, the map $f_i : (X_i, \tau_{X_i}) \rightarrow (Y, \tau_d)$ is a quotient map. Therefore, from **2-d**), the map $h : (Y, \tau_d) \rightarrow (\mathcal{D}_{f_i}, \tau(\mathcal{D}_{f_i})), y \mapsto f_i^{-1}(y)$ must be a homeomorphism. Since (Y, τ_d) is self-similar, according to **2-e**), the decomposition space $\mathcal{D}_{f_i} = \{f_i^{-1}(y) \subset X_i; y \in Y\}, \tau(\mathcal{D}_{f_i}) = \{\mathcal{U} \subset \mathcal{D}_{f_i}; \bigcup \mathcal{U} \in \tau_{X_i}\}$ of (X_i, τ_{X_i}) is also self-similar. The space X is filled with self-similar spaces in the sense that the family $\{f_i^{-1}(y); y \in Y, i = 1, \dots, n\}$ of subsets of X is a cover of X . The self-similar spaces $\mathcal{D}_{f_i}$ and $\mathcal{D}_{f_j}, i \neq j$ are disjoint in the sense that any point $f_i^{-1}(y)$ of $\mathcal{D}_{f_i}$ is not a point of $\mathcal{D}_{f_j}$ (Fig.1).

Finally, we note that in the above discussions we can replace the self-similar space with a compact substance in the materials science such as dendrite (A metric space is called a dendrite provided that it is a connected, locally connected, compact metric space which contains no simple closed curve as its subspace. A space which is homeomorphic to a dendrite is a dendrite. cf. **2-e**.) [1,2] and then, we can obtain a family $\{X_1, \dots, X_n\}$ of subspace X_i whose decomposition space is characterized as a dendrite.

4 Conclusion

Let a topological space (X, τ) be the Λ -product space $(\{0, 1\}^\Lambda, \tau_0^\Lambda)$ of $(\{0, 1\}, \tau_0)$. Here $\text{Card} \Lambda \succ \aleph$ (aleph) and $\tau_0 = \{\{0, 1\}, \{0\}, \{1\}, \emptyset\}$. (X, τ) has an any n -partition $\{X_1, \dots, X_n\}$ where each subspace (X_i, τ_{X_i}) is a non-empty clopen set of (X, τ) and has a self-similar decomposition space. Namely, (X, τ) is filled with the any n mutually disjoint self-similar spaces each of which is a decomposition space of $(X_i, \tau_{X_i}), i \in \{1, \dots, n\}$. It is emphasized that the discussions concerning the self-similarity do not depend on the metrizability of the initial space (X, τ) .

Acknowledgement

The authors are grateful to Dr. Y. Yamashita, Dr. H. Ryo, Prof. Emeritus H. Fukaishi at Kagawa university, and Prof. H. Nagahama at Tohoku university for useful suggestions and encouragements.

References

- [1] A. Kitada, Y. Ogasawara and T. Yamamoto, Chaos, Solitons & Fractals 34 (2007) 1732-1735
- [2] S.B. Nadler Jr., *Continuum theory*, Marcel Dekker (1992) p.165

³As mentioned in **2-f**), there exists 0-dim, perfect, compact, not metrizable T_2 -subspace $(X_{i_0}, \tau_{X_{i_0}})$.